

A LQG extension of the inflationary scenario to the Planck regime

Abhay Ashtekar

Institute for Gravitation and the Cosmos, Penn State

Will summarize the work of many researchers; especially:

Agullo, Barrau, Bojowald, Cailleatau, Campiglia, Corichi, Grain, Kaminski, Lewandowski, Mielczarek, Mena, **Nelson**, Olmedo, Pawlowski, Singh, Sloan, Velhinho ...

AEI Workshop on Quantum Gravity & Fundamental Cosmology, March 5th, 2013.

Setting the Stage

- The Very Early Universe provides a natural arena to test quantum gravity: Guidance for formulating the theoretical framework *and* confronting theories with observations. Challenge: Interface of two communities.
- Standard tools used currently: FLRW solutions to Einstein's equations and quantum field theory of linear perturbations on it, ignoring the back reaction. Checked by self-consistency in the very early universe.
- Challenge to quantum gravity theories: Extend this theory the Planck regime where general relativity breaks down. Do theoretically self consistent extensions exist? Do they pass the current observational tests? Are there new predictions for future observations?
Goal: Probe these issues using Loop Quantum Gravity.
- For concreteness, I will focus on the *inflationary scenario* although the framework is general.

For summary, see: **AA, Agullo & Nelson *PRL* 109, 251301 (2012);**
Viewpoint article, *Physics:Spotlighting Exceptional Research*,
5, 142 (2012.)

Inflationary Paradigm

- Major success: Prediction of inhomogeneities in CMB which serve as seeds for structure formation. Observationally relevant wave numbers in the range $\sim (k_o, 2000k_o)$ (radius of the observable CMB surface $\sim \lambda_o$).
- Rather minimal assumptions:
 1. Some time in its early history, the universe underwent a phase of accelerated expansion during which the Hubble parameter H was nearly constant.
 2. Starting from this phase till the CMB era, the universe is well-described by a FLRW background with linear perturbations. Only matter: inflaton in a suitable potential.
 3. At the onset of this 'slow roll inflationary phase' Fourier modes of quantum fields describing perturbations were in the Bunch-Davies vacuum (at least for co-moving wave numbers in the range $\sim (k_o, 2000k_o)$); and,
 4. Soon after a mode exited the Hubble radius, its quantum fluctuation can be regarded as a classical perturbation and evolved via linearized Einstein's equations.
- Then QFT on FLRW space-times (and classical GR) implies the existence of tiny inhomogeneities in CMB seen by the 7 year WMAP data.
All large scale structure emerged from vacuum fluctuations!

Inflationary Paradigm: Incompleteness

Particle Physics Issues:



- Where from the inflaton? A single inflaton or multi-inflatons? Interactions between inflatons? How are particles/fields of the standard model created during 'reheating' at the end of inflation? ...

Quantum Gravity Issues:

- Big bang singularity also in the inflationary models (Borde, Guth & Vilenkin). Is it resolved by quantum gravity as has been hoped since the 1970's? What is the nature of the quantum space-time that replaces Einstein's continuum in the Planck regime?
- Does the slow-roll inflation used to explain the WMAP data naturally arise from natural initial conditions 'at the Beginning' that replaces the big bang in quantum gravity?
- In classical GR, if we evolve the modes of interest back in time, they become trans-Planckian. Is there a QFT on **quantum** cosmological space-times needed to adequately handle physics at that stage?
- Can one **arrive at** the Bunch-Davies vacuum (at the onset of the WMAP slow roll) from more fundamental considerations?

'Standard' View & its limitations

Could the pre-inflationary dynamics affect the scenario?
Could Planck scale physics play a role?



Inflationary Scenario: Incompleteness

Quantum Gravity Issues:

- Big bang singularity also in the inflationary models (Borde, Guth & Vilenkin). Is it resolved by quantum gravity as has been hoped? Nature of the quantum space-time that replaces Einstein's continuum in the Planck regime?


"One may not assume the validity of field equations at very high density of field and matter and one may not conclude that the beginning of the expansion should be a singularity in the mathematical sense."

A. Einstein, 1945

- In Loop Quantum Gravity, singularities have been resolved in a large number of cosmological models including the (Flat & Closed) FLRW models, Zero and non-zero Λ , Anisotropic Bianchi models, & Gowdy models that have simplest types of inhomogeneities. Mechanism: Quantum Geometry underlying Loop Quantum Gravity. (Bojowald, AA, Lewadowski, Pawłowski, Singh; Wilson-Ewing; D. Brizuela, Martin-Benito, Mena-Marugan).

The Simplest Model


The $k=0$, $\Lambda = 0$ FRW Model coupled to a massless scalar field ϕ .
Instructive because **every** classical solution is singular. **Provides a foundation for more complicated models.**



Classical trajectories

2. Singularity Resolution

$k=0$ LQC with massless scalar field




Absolute value of the physical state $\Psi(v, \phi)$

(AA, Pawłowski, Singh)

Singularity Resolution


k=0 LQC with massless scalar field



Expectations values (and dispersions) of $\hat{V}|_{\phi}$ & classical trajectories.

(AA, Pawłowski, Singh) Gamow's favorite paradigm realized.

Singularity Resolution: $(1/2)m^2\phi^2$ Potential



Expectations values and dispersions of $\hat{V}|_{\phi}$ for a massive inflaton ϕ with phenomenologically preferred parameters (AA, Pawlowski, Singh). The Big Bang is replaced by a Big Bounce.

What is behind singularity resolution?

- In full LQG, we have a mathematically rigorous kinematical framework uniquely selected by the requirement of background independence (Lewandowski, Okolow, Sahlmann, Thiemann; Fleischhack). This descends to LQC in a well defined manner (AA, Campiglia).

- This kinematics is distinct from the Schrödinger representation used in the WDW theory. In particular, the differential operator of the WDW equation, $\partial^2 \Psi_o(v, \phi) / \partial v^2 = \ell_P^2 \hat{H}_\phi \Psi_o(v, \phi)$ fails to be well-defined on the LQC Hilbert space and is naturally replaced by a difference operator:

$$C^+(v) \Psi_o(v+4, \phi) + C^0(v) \Psi_o(v, \phi) + C^-(v) \Psi_o(v-4, \phi) = \ell_P^2 \hat{H}_\phi \Psi_o(v, \phi)$$

- (The Step size is determined by the area gap of Riemannian quantum geometry underlying LQG)

- Singularity Resolution: Not because one ‘jumps over’ the singularity. **Finiteness of physical observables.** The matter density operator $\hat{\rho}$ has an absolute upper bound on the physical Hilbert space (AA, Corichi, Singh):

$$\rho_{\text{sup}} = \sqrt{3}/16\pi^2 \gamma^3 G^2 \hbar \approx 0.41 \rho_{\text{Pl}}!$$

Provides a precise sense in which the singularity is resolved. (Discussion within the consistent histories framework (Criag and Singh)).

Singularity Resolution in LQC: $k=0$

- No unphysical matter. All energy conditions satisfied. But the left side of Einstein's equations modified because of quantum geometry effects (discreteness of eigenvalues of geometric operators.)
- Good agreement with the WDW equation at low curvatures but drastic departures in the Planck regime precisely because the WDW theory ignores quantum geometry (the area gap).
- **Effective Equations:** To compare with the standard Friedmann equation, convenient to do an algebraic manipulation and move the quantum geometry effect to the right side. Then:
$$(\dot{a}/a)^2 = (8\pi G\rho/3)[1 - \rho/\rho_{\text{crit}}] \quad \text{where} \quad \rho_{\text{crit}} \sim 0.41\rho_{\text{Pl}}.$$
Big Bang replaced by a quantum bounce. *Effective equations are surprisingly effective even in the Planck regime.* Simplifies the analysis.
- **Mechanism:** Quantum geometry creates a brand new repulsive force in the Planck regime, neatly encoded in the difference equation. Replaces the big-bang by a quantum bounce.

2. Beyond Homogeneity: Our Approach

- The mainstream strategy in LQG: We do not have a complete quantum gravity theory. Progress has occurred by first truncating the classical theory to the physical problem under consideration and then passing to quantum theory using LQG techniques. Success in the study of quantum horizons, obtaining the graviton propagator, and simple cosmological models.
- For inflation, the sector of physical interest: FLRW background with an inflation ϕ in a suitable potential as matter, together with first order perturbations.
- Our Approach: Use the truncation provided by this cosmological sector. (In numerical simulations, $V(\phi) = (1/2)m^2\phi^2$.) The sector **includes inhomogeneities**, but as perturbations. Thus, quantum fields representing scalar and tensor perturbations now propagate on a **quantum** FLRW geometry. ‘Trans-Planckian’ issues are faced squarely. Caveat: Have to check **self-consistency** of this truncation! Is the back reaction on quantum geometry negligible even in the Planck regime? If so, there we would have a self consistent extension of the inflationary paradigm to the Planck regime. Thus, **two** key issues have to be resolved.

3. Truncated theory


- Truncated Phase Space $\ni \{(v, \phi; \delta h_{ab}(x), \delta \phi(x))$ and their conjugate momenta}

Quantum Theory: Start with $\Psi(v, \phi; \delta h_{ab}(x), \delta \phi(x))$ and proceed to the quantum theory using LQG techniques.

- Test field approximation: $\Psi = \Psi_o(v, \phi) \otimes \psi(\delta h_{ab}, \phi)$, Ψ_o a physical quantum state in the homogeneous sector. Provides QUANTUM background geometry.
- Linearized constraints $\Rightarrow \psi(\delta h_{ab}, \phi) = \psi(T^{(1)}, T^{(2)}, \mathcal{R}; \phi)$, where $T^{(1)}, T^{(2)}$ are the tensor modes and \mathcal{R} the scalar mode. In the Planck regime of interest, ϕ serves as the ‘internal/relational time’. ψ propagates on the quantum geometry determined by Ψ_o .
- Idea: Choose $\Psi_o(v, \phi)$ to be sharply peaked at an effective LQC solution g_{ab}^o . Such ‘coherent states’ exist. First question: Does the required inflationary phase occur generically in such quantum geometries Ψ_o ?

Quantum FLRW Background Geometry Ψ_o

- Let us start with generic data at the bounce in the effective theory and evolve. Will the solution enter slow roll at energy scale $\rho \approx 7.32 \times 10^{-12} m_{\text{Pl}}^4$ determined from the 7 year WMAP data ? Note: 11 orders of magnitude from the bounce to the onset of the desired slow roll!
- Answer: **YES**. In LQC, $|\phi_B| \in (0, 7.47 \times 10^5)$. If $\phi_B \geq 0.93$, the data evolves to a solution that encounters the slow roll **compatible with the 7 year WMAP data** sometime in the future. In this sense, ‘almost every’ initial data at the bounce evolves to a solution that encounters the desired slow roll sometime in the future. (AA & Sloan, Corichi & Karami)
- Result stronger than the ‘attractor’ idea because it refers to the slow roll compatible with WMAP.
- Hence, for the background quantum geometry, we can choose a ‘coherent’ state Ψ_o sharply peaked at an effective trajectory with $\phi_B > 0.93$ and evolve using LQC. **WMAP slow roll phase ensured!**



Inflationary Paradigm: Incompleteness

Quantum Gravity Issues:

- Big bang singularity also in the inflationary models (Borde, Guth & Vilenkin). Is it resolved by quantum gravity as had been long hoped? What is the nature of the quantum space-time that replaces Einstein's continuum in the Planck regime?
- In the systematic evolution from the Planck regime in the more complete theory, does a slow roll phase compatible with the WMAP data arise **generically** or is an enormous **fine tuning** needed?
- In classical GR, if we evolve the modes of interest back in time, they become trans-Planckian. Is there a QFT on **quantum** cosmological space-times needed to adequately handle physics at that stage? **Yes!**
The basic framework was developed in 2009 (AA, Lewandowski, Kaminski).
Can it be further developed to obtain well-defined stress energy operators in this new theory to examine if the back reaction is negligible, i.e. **justify the truncation procedure** even in the Planck regime?
- Can one **arrive at** the Bunch-Davies vacuum (at the onset of the WMAP slow roll) from more fundamental considerations?

4. Extracting Physics

First, thanks to the background **quantum** geometry, trans-Planckian modes pose no problem, **provided** the test field approximation holds: $\rho_{\text{Pert}} \ll \rho_{\text{BG}}$ all the way from the bounce to the onset of slow roll.

- Second, surprisingly, truncated dynamics of $\hat{T}^{(1)}, \hat{T}^{(2)}, \hat{\mathcal{R}}$ on the **quantum** geometry of Ψ_o is mathematically equivalent to that of $\hat{T}^{(1)}, \hat{T}^{(2)}, \hat{\mathcal{R}}$ as quantum fields on a smooth space-time with a ‘**dressed**’ effective, **c-number metric** \bar{g}_{ab} (whose coefficients depend on \hbar):

$$\bar{g}_{ab} dx^a dx^b = \bar{a}^2 (-d\bar{\eta}^2 + d\vec{x}^2)$$

with

$$d\bar{\eta} = \langle \hat{H}_o^{-1/2} \rangle [\langle \hat{H}_o^{-1/2} \hat{a}^4 \hat{H}_o^{-1/2} \rangle]^{1/2} d\phi; \quad \bar{a}^4 = (\langle \hat{H}_o^{-1/2} \hat{a}^4 \hat{H}_o^{-1/2} \rangle) / \langle \hat{H}_o^{-1} \rangle$$

where H_o is the Hamiltonian governing dynamics of Ψ_o . **Analogy with light propagating in a medium.**

- Because of this, the mathematical machinery of adiabatic states, regularization and renormalization of the Hamiltonian can be **lifted to** the QFT on cosmological QSTs under consideration. Result: Full mathematical control on **dynamics** for computation of the CMB power spectrum, and spectral indices starting from the bounce.

Initial conditions on ψ

- Ψ_o : **special** because assumed to be a ‘coherent state’ at the bounce. But peaked at a **generic** effective trajectory.
- ψ : Hilbert space \mathcal{H} of perturbations on the quantum geometry Ψ_o is spanned by 4th adiabatic order states on the smooth Friedmann metric \bar{g}_{ab} . Excellent control.
- Initial conditions: Since Ψ_o (and hence \bar{g}_{ab}) is homogeneous and isotropic, demand that: (i) $\psi \in \mathcal{H}$ also invariant under translations and rotations; (ii) Back reaction of the perturbation ψ on the background Ψ_o can be ignored at the bounce; and (iii) ‘Fundamental uncertainties minimized at the bounce’.
- Intuitive Physical Meaning: Demanding initial **quantum** homogeneity and isotropy. Heuristic justification: Because of inflation, the observable universe has size of $\leq 10\ell_{\text{Pl}}$ at the bounce. The **repulsive force of quantum geometry** dilutes all inhomogeneities at this scale. So universe is as homogeneous and isotropic as the uncertainty principle allows it to be! Uniqueness issue still being explored. So far the emphasis is on **existence** of initial conditions that provide a satisfactory extension of the inflationary scenario.

Key Questions: Agullo's talk

1. Does the back-reaction remain negligible as ψ evolves all the way to the onset of the slow roll compatible with WMAP (so that our truncation strategy is justified by self-consistency)?

Brief Answer: YES.

2. At the end of the WMAP compatible slow roll, do we recover the observed power spectrum: $\Delta_{\mathcal{R}}^2(k, t_{k^*}) \approx \frac{H^2(t_{k^*})}{\pi m_{\text{Pl}}^2 \epsilon(t_{k^*})}$? (t_{k^*} is the time the reference mode $k^* \approx 8.58k_o$ exits the Hubble horizon during slow roll)

Brief Answer: YES. Thus, there is a self-consistent quantum gravity completion of the inflationary paradigm.

3. Does $\psi(T_{\bar{k}}^{(1)}, T_{\bar{k}}^{(2)}, \mathcal{R}_{\bar{k}}; \phi_B)$ evolve to a state which is indistinguishable from the Bunch Davies vacuum at the onset of slow roll or are there deviations with observable consequences for more refined future observations (e.g. non-Gaussianities in the bispectrum)? (Agullo & Shandera; Ganc & Komatsu; Schmidt & Hui, ...)

Brief Answer: There **ARE** deviations for a small window in the parameter space provided by ϕ_B .

5. Summary: Framework

- The early universe provides an ideal setting to test quantum gravity ideas. Can one obtain a quantum gravity extension of cosmological scenarios to the Planck regime? We focused on inflationary paradigm because it has been extremely successful with structure formation.

The standard theory can be extended using LQG:

- *Background geometry*: Singularity Resolution and precise quantum geometry for the Planck regime. ✓
- *Perturbations*: Since they propagate on quantum geometry, using QFT on cosmological quantum geometries (AA, Lewandowski, Kaminski), trans-Planckian issues can be handled systematically provided the test field approximation holds. Analyzed in detail using the renormalized stress-energy of $\hat{T}^{(1)}$, $\hat{T}^{(2)}$, $\hat{\mathcal{R}}$ on the quantum geometry of Ψ_o . Detailed numerics show that the approximation does hold in most of the parameter space. (Agullo, AA, Nelson). ✓
Detailed calculations in the inflationary paradigm but framework is much more general.

Summary: Implications for Inflation

- **Extension:** For most of the parameter space, modes of observational interest are essentially in the Bunch Davies vacuum at the onset of the WMAP slow roll \Rightarrow Predictions of the standard inflationary scenario for the power spectra, spectral indices & ratio of tensor to scalar modes are recovered starting from Planck era. (Agullo, AA, Nelson) ✓
- **Non-Gaussianity:** There is a small window in the parameter space, for which at the onset of inflation ψ has excitations over the Bunch-Davies vacuum. These give rise to specific 3-point functions ('bi-spectrum') which are important for the 'halo bias'. Could be observed in principle: Link between observations and the initial state! A window to probe the Planck era around the LQC bounce. (Agullo, AA, Nelson, Shandera, Ganc, Komatsu) ✓
- **Note:** LQG does **not** imply that inflation **must** have occurred because it does not address particle physics issues. The analysis simply **assumes** that there is an inflaton with a suitable potential. But it does show concretely that many of the standard criticisms (e.g. due to Brandenberger) of inflation can be addressed in LQG by facing the Planck regime squarely. The framework is general; can be applied to other viable scenarios.

Main References for this talk

- For a summary, see:

AA, Agullo & Nelson PRL 109, 251301 (2012);

- More complete references:

AA, Agullo & Nelson, PRD 87, 043507 (2013); 1302.0254

AA Sloan GRG (2011), PLB (2009);

AA, Corichi & Singh PRD (2008); AA, Pawłowski, Singh, PRL & PRD (2006).

- Future Observations:

Agullo & Parker PRD & GRG (2011); Agullo & Shandera JCAP (2012);
Ganc & Koamtzu PRD (2012).

- A recent detailed Review of Loop Quantum Cosmology

AA & Singh, CQG (2011).

Supplementary Material

The slides that follow represent supplementary material, not included in the main talk. They address some general questions and provide computer simulations showing that the program discussed in this talk has been completed. Details on this completion, and on phenomenological and observational issues can be found in Ivan Agullo's talk.

Merits and Limitations of QC

One's first reaction to Quantum Cosmology is often: Symmetry reduction gives only toy models! Full theory much richer and much more complicated.

But examples can be powerful.

- Full QED versus Dirac's hydrogen atom.
- Singularity Theorems versus first discoveries in simple models.
- BKL behavior: homogeneous Bianchi models.

Do *not* imply that behavior found in examples is necessarily generic. Rather, they can reveal important aspects of the full theory and should not be dismissed a priori.

One can work one's way up by considering more and more complicated cases. (e.g. recent work of the Madrid group on Gowdy models which have infinite degrees of freedom). At each step, models provide important physical checks well beyond formal mathematics. Can have strong lessons for the full theory.

Remaining Key Questions

Return to our quantum gravity extension of the inflationary paradigm.

1. Does the back-reaction remain negligible as ψ evolves all the way to the onset of the slow roll compatible with WMAP (so that our truncation strategy is justified by self-consistency)?

Answer: **YES**

2. At the end of the WMAP compatible slow roll, do we recover the observed power spectrum: $\Delta_{\mathcal{R}}^2(k, t_{k^*}) \approx \frac{H^2(t_{k^*})}{\pi m_{\text{Pl}}^2 \epsilon(t_{k^*})}$? (t_{k^*} is the time the reference mode $k^* \approx 8.58k_o$ exits the Hubble horizon during slow roll)

Answer: **YES provided $\phi_B \geq 1.14m_{\text{Pl}}$** . Thus, we have arrived at a **quantum gravity completion** of the inflationary paradigm.


3. Does $\psi(T_{\bar{k}}^{(1)}, T_{\bar{k}}^{(2)}, \mathcal{R}_{\bar{k}}; \phi_B)$ evolve to a state which is indistinguishable from the Bunch Davies vacuum at the onset of slow roll or are there deviations with observable consequences for more refined future observations (e.g. non-Gaussianities in the bispectrum)?

(Agullo & Shandera; Ganc & Komatsu)

Answer: There **ARE** deviations if ϕ_B lies in a small window just after $\phi_B = 1.14m_{\text{Pl}}$.


Self-consistency of Truncation: $\rho_{\text{Pert}}/\rho_{\text{BG}}$ vs time.

Question 1: We have explicit numerical simulation showing that the truncation strategy is self-consistent.



Renormalized energy density in ψ is negligible compared to that in Ψ_o all the way from the bounce to the onset of slow roll. Here $\phi_B = 1.15m_{\text{Pl}}$.

The Scalar Power spectrum: Ratio ($\mathcal{P}_{\text{LQG}}/\mathcal{P}_{\text{BD}}$)




Question 2:

Ratio of the LQG and the standard BD power spectrum for the scalar mode. Blue: Raw data points. Red: Average. LQG prediction is within observational errors for $\phi_B \geq 1.14m_{\text{Pl}}$.

For $\phi_B = 1.2m_{\text{Pl}}$, WMAP $k_{\text{min}} = 9m_{\text{Pl}}$. Complete agreement with BD vacuum for $\phi_B \geq 1.2m_{\text{Pl}}$. For $\phi_B < 1.2m_{\text{Pl}}$: Certain non-Gaussianities for future observations.

The LQC Tensor Power spectrum



Question 2: Tensor Modes:

Predicted power spectra for the tensor mode. Black: Average. Red: Raw data points.

Understanding the Power Spectrum

Question 3: Only modes with $\lambda > R_{\text{curv}}$, the curvature radius, in the pre-inflationary era are excited and populated at the onset of inflation. Can occur in a narrow window for $\phi_B \leq 1.2m_{\text{Pl}}$.

